

## FUSE Observations of $\eta$ Carinae

R. C. Iping, G. Sonneborn, T. R. Gull

*NASA's GSFC, Code 681, Greenbelt, MD 20771*

**Abstract.**  $\eta$  Carinae was observed by FUSE through the LWRS ( $30'' \times 30''$ ) and HIRS ( $1.25'' \times 20''$ ) apertures in March and April 2004. There are significant differences between the two spectra. About half of the LWRS flux appears to be due to two B-type stars near the edge of the LWRS aperture,  $14''$  from  $\eta$  Car. The HIRS spectrum (LiF1 channel) therefore reveals the intrinsic FUV spectrum of  $\eta$  Car without this stellar contamination. The HIRS spectrum contains strong interstellar  $H_2$  having high rotational excitation (up to  $J = 8$ ). Most of the atomic species with prominent ISM features (C II, Fe II, Ar I, P II, etc) also have strong blue-shifted absorption to  $v \sim -580$  km/s that is associated with expanding debris from the 1840 eruption.

### 1. $\eta$ Carinae Background

The luminous blue variable (LBV)  $\eta$  Carinae (HD 93308), a highly evolved, massive and luminous star, shows strong evidence for a binary companion with a 5.5 year periodicity in X-rays, optical and UV (Daminelli 1996, van Genderen et al. 2003, Whitelock et al. 2004, Corcoran et al. 2001). Most of its luminosity is emitted in the infra-red, but there is flux and variability at all wavelengths from hard X-rays to radio. The observed double-lobed structure consists of ejecta from the massive outburst in 1840. The orbital parameters of  $\eta$  Car are not well constrained, although extensive modelling is on its way (Corcoran et al. 2001; Pittard & Corcoran 2002; Corcoran et al. 2005, in preparation).

### 2. The FUSE Spectrum

In this paper we report on observations for our FUSE Cycle 4 GI program D007 obtained in early 2004.  $\eta$  Car was observed through the LWRS aperture ( $30'' \times 30''$ ) on Mar. 29-20 (p.a.  $\sim 120^\circ$ ) and the HIRS aperture ( $1''.25 \times 20''$ ) on Apr. 9-11 (p.a.  $\sim 131^\circ$ ). The exposure times were 41 ksec (LWRS) and 83 ksec (HIRS), yielding a high S/N spectrum, part of which is shown in Figure 1. The observed LWRS flux at  $1040 \text{ \AA}$  is  $\sim 3 \times 10^{-12} \text{ erg cm}^{-2} \text{ sec}^{-1} \text{ \AA}^{-1}$ . The flux level declines toward the Lyman limit where converging  $H_2$  and H I features completely blanket the spectrum. In the HIRS aperture, after correcting for the 60% point-source throughput, the  $1040 \text{ \AA}$  flux is only  $1.5 \times 10^{-12}$ . The flux difference appears to be due to two  $10^{th}$ -magnitude B-type stars, Tr16-64 and Tr16-65, that are  $14''$  from  $\eta$  Car and fall just inside the LWRS aperture. The inferred FUV flux for these stars ( $\sim 1.5 \times 10^{-12}$ ) is consistent with their brightness in HST/ACS  $2200 \text{ \AA}$   $2500 \text{ \AA}$  and  $3300 \text{ \AA}$  images obtained in 2003 (Smith




Figure 1. The LiF1a spectrum of  $\eta$  Car from 990 Å to 1070 Å. The thick line of the HIRS aperture spectrum; the thin line is LWRS. The HIRS flux was scaled up by 1.67X to correct for the point source throughput (60%). The  $H_2$  Lyman and Werner ground state transitions are shown, respectively, by short- and long-dashed lines and are identified by the rotational branch and quantum number (e.g. R3) above each line and by the vibrational quantum numbers (e.g. 0-8) on the left-hand side of each panel. The principal atomic transitions and the R0 lines of HD are also identified. All transitions are marked at their laboratory wavelengths.




Figure 2. 1130-1140Å portion of  $\eta$  Car LWRS spectrum (thin) and HIRS spectrum (thick). Note the large difference in Fe II & N I profiles from 1133-1133.5Å. This is due to two B-stars just within the LWRS that are excluded by the HIRS aperture. The horizontal lines indicate 0 to  $-585 \text{ km s}^{-1}$  and mark the absorption by circumstellar debris from the eruptions in the 1800s. The absorption profiles of C I 1138.56, 1139.79Å also indicate ejecta absorption.

et al. 2004). The throughput-corrected HIRS FUV flux from  $\eta$  Car at 1150-1180Å is 10X higher than that observed in HST/STIS E140M spectra ( $0''.2 \times 0''.2$  aperture) at the same wavelengths, demonstrating that there is a considerable circumstellar/scattered component to the observed FUV flux.

Numerous broad features ( $\sim 1 \text{ Å}$ ) in Fig. 1 and 2 are identified as absorption by a forest of high-velocity narrow lines formed in the expanding circumstellar debris of  $\eta$  Car. These features span a heliocentric velocity range of  $-140$  to  $-580 \text{ km s}^{-1}$  and are seen prominently in low-ionization ground-state transitions (e.g. N I, Fe II, P II, Ar I, C I, C III 1175-6, and several unidentified features (see Fig. 2). The high-velocity components are well-resolved in HST/STIS echelle spectra (Nielsen et al. 2005, ApJS in press).

**Acknowledgments.** These observations were obtained through the FUSE GI program D007. We are indebted to the positive support of B-G Andersson and the FUSE mission planning team.

## References

- Corcoran, M.F., Ishibashi, K., Swank, J. H., & Petre, R. 2001, ApJ, 547, 1034
- Daminelli, A. 1996, ApJ, 460, L49
- van Genderen, A., Sterken, C., Allen, W. H., & Lillier, W. 2003, A & A, 412, L25
- Pittard, J.M., & Corcoran, M.F. 2002, A & A, 383, 636
- Smith, N., et al. 2004, ApJ, 605, 405
- Whitelock, P., Feast, M.W., Marang, F., & Breedt, E. 2004, MNRAS, 352, 447